






ICD-10 CM CODE STRUCTURE								>>		
						_				
IC	D-9-C	M								
		7	2	0						
		/	2	7		5				
	- 40	~								
IC	D-10-0	JM								
	ΝЛ	7	O		4	2	1			
	IVI		7		U					



























ucf-rec.org


Varicose veins associated with pregnancy


6 UNIVERSITY OF CENTRAL FLORIDA




























UCF REGIONAL EXTENSION CENTER **7TH CHARACTER FOR FETUS IDENTIFICATION** I.C.15.a.6 • Where applicable, a 7th character is to be assigned for certain categories to identify the fetus for which the complication code • Assign 7th character "0" For single gestations When the documentation is insufficient to determine the fetus affected and it is not possible to obtain clarification When it is not possible to clinically determine which fetus is affected UCF REGIONAL EXTENSION CENTER **OUTPATIENT PRENATAL VISITS** I.C.15.b.1 and 2: Outpatient prenatal visits • For routine outpatient prenatal visits when no complications are present, a code from category Z34 should be used as the first-listed diagnosis. These codes should not be used in conjunction with chapter 15 codes. • For routine prenatal outpatient visits for patients with high-risk pregnancies, a code from category O09 should be used as the first-listed diagnosis. Secondary chapter 15 codes may be used in conjunction with these codes if appropriate. ucf-rec.org UCF REGIONAL EXTENSION CENTER **DELIVERY EPISODES** I.C.15.b.4 and 5 When a delivery occurs, the principal diagnosis should correspond to the main circumstances or complication of the delivery If the patient was admitted with a condition that resulted in the performance of a C-section, that condition should be selected as the principal diagnosis. If the reason for the encounter was unrelated to the condition resulting in the cesarean delivery, the condition related to the reason for the encounter should be selected as the principal diagnosis. A code from category Z37 is to be used on the *maternal* record when a delivery has occurred. This code is not used on subsequent records or on the newborn record

